

Proportion of and factors associated with Thyroid Dysfunction among individuals coming to a tertiary care facility in Kabul,Afghanistan



Mohammad Naeem, Ambreen Gowani and Ahmed Nasir

Clinical Pathologist Afghanistan

Abstract

Background: The thyroid gland, located just below the vocal cord on each side of and anterior to the trachea, is one of the main endocrine glands. Its normal weight is 15 to 20 grams in adults. The thyroid secretes two most important hormones, thyroxine and triiodothyronine, usually called T4 and T3, respectively. These hormones greatly increase the metabolic rate of the body. In addition to T3 and T4, the thyroid gland secrets calcitonin as well which is a signifi67890c-ant hormone for calcium metabolism.

Objective: The aim of this study is to estimate the proportion of and to identify factors associated with thyroid dysfunction among individuals coming to a tertiary care facility in Kabul, Afghanistan.

Material and Methods: An analytical cross-sectional study design was conducted from July to Sep 2018. Blood samples were obtained, serum TSH levels were measured, and the patients were divided into three diagnostic categories according to their serum TSH concentrations. 1) Hypothyroidism 2) Hyperthyroidism 3) Normal thyroid

Results: A total of 127 individuals were part of the sample for the final analysis. Majority study participants (77%) were females. A large number of the members (92%) did not have family history of thyroid brokenness and larger part of the female members, (85%) were not pregnant within the final two a long time. Moreover, a 98% members were non-smokers.

Conclusion: The findings of the current study showed a high prevalence of thyroid dysfunctions in individuals coming to FMIC for thyroid functions tests. The findings also indicated that aging and smoking are the factors associated with thyroid dysfunctions. Further studies are needed to find out the prevalence of and factors associated with thyroid dysfunctions.



Biography

Mohammad Naeem has completed his M.D degree at the age of 28 years from Shaikh Zayed University, Khost Afghanistan. He is the Head of Clinical Laboratory in Noor Clinic here in Khost Afghanistan

10th Annual Congress On Medicinal Chemistry and Drug Design | Amerstdam | Netherlands | March 25, 26, 2020

Citation: Mohammad Naeem, *Proportion of and factors associated with Thyroid Dysfunction among individuals coming to a tertiary care facility in Kabul, Afghanistan* | 10th Annual Congress on Medicinal Chemistry and Drug Design | Mar 25-26, 2020 | Amerstdam, The Netherlands